

Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin Govind Ballabh Pant University of Agriculture & Technology



Udham Singh Nagar, Pantnagar, Uttarakhand

Agromet Advisory Bulletin

Date: 03-03-2023

Weather Forecast of District UDAMSINGH-NAGAR(Uttarakhand) Issued On: 2023-03-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-03-04	2023-03-05	2023-03-06	2023-03-07	2023-03-08
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	31.0	32.0	32.0	32.0	32.0
Tmin(°C)	11.0	11.0	10.0	10.0	10.0
RH-I(%)	50	40	30	30	25
RH-II(%)	15	15	10	15	15
Wind Speed(kmph)	15.0	12.0	13.0	10.0	10.0
Wind Direction(Degree)	300	300	110	300	30
Cloud Cover(Octa)	0	2	0	1	1

Weather Summary/Alert:

In the last seven days (24 Feb -02 March, 2023), there was 0.0 mm rain and the sky was clear to p-cloudy. The maximum & minimum temperatures ranged between 27.5 to 31.0°C and 8.8 to 12.4°C respectively. Relative humidity ranges from 79 to 92 % at 0712 hrs and 28 to 47 % at 1412 hrs. The wind was mainly from the WSW & WNW with a speed of 0.5 to 5.4 km/hour. In the coming five days, weather will remain clear. Maximum & minimum temperature may range from 31.0 to 32.0 deg C & 10.0 to 11.0 deg C respectively. Wind will blow with a speed of 10.0-15.0 km/hr from the E, NW, SE & ESE direction.

General Advisory:

Farmers are advised to download the "Meghdoot App" to get past week weather, weather forecast, and agromet advisory and "Damini App" to get lightning information. Meghdoot and Damini Apps can be downloaded from the google play store (Android users) and App Center (IOS users).

SMS Advisory:

In the coming five days, weather will remain clear. Farmers are advised to irrigate the crops/orchards to reduce the impact of high temperature.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
	On the occurrence of aphid in wheat, spray the solution of Thiomethaxam 25 WSG @ 100
	ml/ha or Imidacloprid 17.8 SL @ 140 ml/ha in 500-600 liters of water. Do second spray at

Crop(Varieties)	Crop Specific Advisory
	Crop(Varieties) Crop Specific Advisory an interval of 15 days. Spray should be done in the morning or evening hours when there is no rain or strong wind. In the timely sown wheat crop, irrigate should be applied at flowering stage based on weather forecast.
	On the highly occurrence of aphid in mustard, Methyl-o-demeton 25 ec @ 1.5 lit/ha or malathion 50 ec @ 2.0 lit/ha should be sprayed by keeping in view the weather forecast.
SUGARCANE	Sowing of spring season sugarcane crop should be completed till 15 March.
LENTIL	Irrigation should be arranged as per requirement in lentil crop at grain filling stage.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory	
MANGO	If Mango mealy bug appeared in the mango trees then cut and separate it and put it in the pit, cover with soil. Last year Gumba baur or Mango mal formation and leaves which are highly affected by powdery mildew should be destroyed by plucking.	
TOMATO	In case of late blight, spray Mancozeb 64% + cymoxanil 8% WP @ 2.5gm per litre by keeping in view the weather forecast.	
BOTTLE GOURD	Field preparation should be done for sowing bottle gourd (lauki), bitter gourd (Karela), ridged gourd (Torai) and pumpkin (Kashiphal) crop. Compost should be used as nutrients for Cucumber class crop.	
VEGETABLE PEA	When lower leaves of pea are turning yellow and plants are drying, then roots of pea crop should be irrigated with the solution of carbendazim @ 1 g/ lit of water	
CABBAGE	BBAGE Top dressing of Urea should be done in Cabbage and Cauliflower crop after weeding and hoeing and moisture should be maintained in the field.	

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
BUFFALO	Disbudding or destroy the horn buds of calves of cow/buffalo at the age of one week.
ICTCJA I	Goats and sheep should be vaccinated by sheep/goat pox to protect them from chickenpox.