

Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Govind Ballabh Pant University of Agriculture & Technology Udham Singh Nagar,Pantnagar, Uttarakhand



Agromet Advisory Bulletin

Date : 14-03-2023

Weather Forecast of District UDAMSINGH-NAGAR(Uttarakhand) Issued On : 2023-03-14(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-03-15	2023-03-16	2023-03-17	2023-03-18	2023-03-19
Rainfall(mm)	0	0	0	3	0
Tmax(°C)	32.0	31.9	31.7	29.6	27.8
Tmin(°C)	10.8	12.1	12.2	10.9	11.2
RH-I(%)	41	30	40	53	49
RH-II(%)	17	13	13	18	24
Wind Speed(kmph)	10.6	10.4	9.6	13.8	10.7
Wind Direction(Degree)	34	56	60	45	72
Cloud Cover(Octa)	1	4	7	7	7

Weather Summary/Alert:

In the last seven days (7 - 13 March, 2022), there was 0.0 mm rain and the sky was Clear to cloudy. The maximum & minimum temperatures ranged between 28.5 to 31.5°C and 11.5 to 15.6°C respectively. Relative humidity ranges from 77 to 93 % at 0712 hrs and 26 to 52 % at 1412 hrs. The wind was mainly from the WNW & WSW with a speed of 0.2 to 7.0 km/hour. In the coming five days, light rain may occur on 17 March and weather will remain clear in the remaining days. Maximum & minimum temperature may range from 27.5 to 32.0 deg C & 10.5 to 12.5 deg C respectively. Wind will blow with a speed of 9.0-14.0 km/hr from the WNW & ESE direction.

General Advisory:

Farmers are advised to download the "Meghdoot App" to get past week weather, weather forecast, and agromet advisory and "Damini App" to get lightning information. Meghdoot and Damini Apps can be downloaded from the google play store (Android users) and App Center (IOS users)

SMS Advisory:

In the coming five days, light rain may occur on 17 March and weather will remain clear in the remaining days

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
WHEAT	Necessary irrigation should be given in the wheat crop as needed. Remove weeds like Gulli Danda and jalgi jai (wild oats) and destroy them. On the occurrence of aphid in wheat, spray the solution of Thiomethaxam 25 WSG @ 100 ml/ha or Imidacloprid 17.8 SL @ 140 ml/ha in 500-600 liters of water. Do second spray at an interval of 15 days. Spray should be done in the morning or evening hours when there is no rain or strong wind.
MUSTARD	Harvest the mature crop of black mustard (rai) and store the grain after drying.
CHICK PEA	Light irrigation should be given in the chickpea crop at the time of pod formation. To control the blight disease in gram, Mancozeb 75% WP @ 2.5g/lit should be sprayed.
SUGARCANE Irrigate the crop sown in the month of February and remove weeds by hoeing after Complete the sowing of spring sugarcane crop. One line of urd, moong or cowpear sown as intercrops between two rows of sugarcane. Irrigate the winter crop of sug per requirement and do top dressing of urea as per recommendation.	
LENTIL	To control the blight disease in lentil, Mancozeb 75% WP @ 2.5g/lit should be sprayed

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory		
POTATO	Complete the digging of potatoes. Potato rot starts with late digging.		
TOMATO	On the occurrence of fruit borer in the Tomato, fruit should be used only after 3 days and 5 days of spraying of Chlorantraniliprole 18.5 SC @ 150ml/Ha and Indoxacarb 14.5 SC @ 500ml/Ha respectively		
ONION	Weeding, hoeing and irrigating should be done in onion crop as per the requirement.		

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory	
	Sitting place of animals should be leveled so that their productive capacity is not affected and special care should be given to the new born animals.	