

# Gramin Krishi Mausam Sewa

# District Level Agromet Advisory Bulletin Govind Ballabh Pant University of Agriculture & Technology



Udham Singh Nagar, Pantnagar, Uttarakhand

# **Agromet Advisory Bulletin**

Date: 23-12-2022

Weather Forecast of District UDAMSINGH-NAGAR(Uttarakhand) Issued On: 2022-12-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-24	2022-12-25	2022-12-26	2022-12-27	2022-12-28
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	26.0	24.0	21.0	22.0	22.0
Tmin(°C)	3.0	4.0	5.0	4.0	4.0
RH-I(%)	90	90	90	90	90
RH-II(%)	30	30	30	30	30
Wind Speed(kmph)	6.0	6.0	6.0	6.0	6.0
Wind Direction(Degree)	320	320	320	330	330
Cloud Cover(Octa)	0	1	4	1	0

## Weather Summary/Alert:

In the last seven days (16 - 22 Dec, 2022), there was 0.0 mm rain and the sky was clear. The maximum & minimum temperatures ranged between 20.0 to 25.0°C and 6.0 to 9.1°C respectively. Relative humidity ranges from 84 to 97 % at 0712 hrs and 44 to 67 % at 1412 hrs. The wind was mainly from the WNW & ENE with a speed of 0.5 to 2.3 km/hour. In the coming five days, weather will remain clear. Maximum & minimum temperature may range from 21.0-26.0 deg C & 3.0 - 5.0 deg C respectively. Wind will blow with a speed of 6.0 km/hr from the NW direction. As per ERF, rainfall will be below normal, maximum & minimum temperature will be moderately above normal respectively in the state from 28 Dec to 3 Jan.

#### **General Advisory:**

IMD Weather forecast and Agrometeorological advisory of your location is now available on Meghdoot mobile app in English and regional languages. Kindly download from following link Android: https://play.google.com/store/apps/details?id=com.aas.meghdoot iOS: https://apps.apple.com/in/app/meghdoot/id1474048155

## **SMS Advisory:**

In the coming five days, weather will remain clear. Farmers are advised to irrigate the crop as per requirement.

## **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory		
WHEAT	In timely sown wheat, first irrigation should be given after 20-25 days of sowing. After 3-4 days of first irrigation, remaining dose of Nitrogen should be applied as top dressing in the afternoon. To control the weed in the wheat, first and second weeding and hoeing should be done after 25-30 days and 45-50 days of sowing respectively. Sowing of late sown variety of wheat should be done till 25 December. In case of late sowing, decrease in temperature has an adverse effect on germination and growth of wheat crop. So it is advise that seed rate should be 25% more. Sowing should be done at a distasnce of 15-18 cm in lines and at a depth of 3-4 cm.		
MUSTARD	Light irrigation should be given at the time of grain filling in late sown crops of toria and yellow mustard. To control the leaf spot disease in mustard, solution of mancozeb @ 2.5 g/liter of water should be sprayed. Harvesting of yellow mustard & toria, sown in the month of September should be done on maturity.		
BARLEY	Late sowing variety of Barley like-Jyoti, Preeti, Manjula, Jagrati should be selected. Seed rate should be 100-110Kg seed/Ha. Sowing of seeds should be done at 18-20cm in the lines and comple the sowing by second fortnight of December.		
SUGARCANE	Weeding and hoeing should be done in the sugarcane crop and remaining dose of nitrogen should be applied on time.		

# **Horticulture Specific Advisory:**

Horticulture(Varieties)	Horticulture Specific Advisory		
TOMATO	On the occurrence of yellowish brown spot in tomato, mancozeb 2.5-3g/lit of water should be sprayed.		
BRINJAL	On the withering of brinjal, solution of carbendazim @1 g/ lit of water should be sprayed. If whole plant is drying then root should be irrigated with the same solution.		
CHILLI	On the shrinkage or getting piebald of upper leaves of chilly and tomato, infected plants should be removed and destroyed. To control the disease carried insects any sarvangi insecticide should be sprayed.		
VEGETABLE PEA	When lower leaves of pea are turning yellow and plants are drying, then roots of crop should be irrigated with the solution of carbendazim @ 1 g/ lit of water.		
MANGO	To stop the Gujjiya keet for climbing on a tree, the stem of the tree should be covered by fine soil layer up to 40 cm height from the ground in the last week of this month and wrap up with 25 cm broad strip of 400 gazes thick polythene and tie with the help of sutali. Along with this grease must be applied in the lower portion.		

# Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
BUFFALO	To save the animal from cold, increase the amount of oil and Jaggery (Gur) in the food of animals. Caraway (Ajwain) and gur also be given to the animals. To save the animals from cold, proper arrangement of animal shed should be done. Vaccination should be given to the animals to save them from Rinderpest disease (Sheetla rog).