

Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Govind Ballabh Pant University of Agriculture & Technology Udham Singh Nagar,Pantnagar, Uttarakhand



# **Agromet Advisory Bulletin**

Date : 28-01-2022

Weather Forecast of NAINITAL(Uttarakhand) Issued On : 2022-01-28(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-01-29	2022-01-30	2022-01-31	2022-02-01	2022-02-02
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	10.0	12.0	12.0	13.0	12.0
Tmin(°C)	0.0	1.0	2.0	1.0	2.0
RH-I(%)	65	55	50	45	45
RH-II(%)	80	80	75	75	65
Wind Speed(kmph)	6.0	6.0	8.0	6.0	6.0
Wind Direction(Degree)	320	320	320	320	320
Cloud Cover(Octa)	1	1	0	0	1

#### Weather Summary/Alert:

In the coming five days weather will remain clear. Maximum & minimum temperature may range from 10.0 to 13.0 deg C & 0.0 to 2.0 deg C respectively. Wind will blow with a speed of 6.0-8.0 km/hr from the NW direction. There is a warning of likely occurrence of ground frost at isolated places in hills from 28 to 30 Jan. As per ERF, the rainfall will be above normal, maximum & minimum temperature will be below normal in the state from 2 to 8 February.

#### **General Advisory:**

NDVI & SPI map received from the India Meteorological Department dated indicated that the NDVI ranged between 0.2-0.5 for Nainital i.e. agriculture vigor is moderate to good in the district. SPI map indicates the extremely wet conditions in the district during 30 Dec 2021 to 26 Jan 2022. Farmers are advised to download the "Meghdoot App" to get past week weather, weather forecast, and agromet advisory and "Damini App" to get lightning information. Meghdoot and Damini Apps can be downloaded from the google play store (Android users) and App Center (IOS users). Weather will remain clear in the coming days, so farmers can apply the irrigation & fertilizer in the field.

#### **SMS Advisory:**

In the coming five days weather will remain clear. Ground frost may likely occur at isolated places in hills, so the farmers can protect the crops from frost by applying sprinkle irrigation.

#### **Crop Specific Advisory:**

Crop(Varieties) Crop Specific Advisory
--

District Advisory

<b>Crop(Varieties)</b>	Crop Specific Advisory
WHEAT	On the occurrence of rust in wheat crop, solution of Propiconazole 25 EC @ 1ml/liter of water should be sprayed by keeping in view the weather forecast.
MUSTARD	So, on the occurrence of Sclerotinia rot in mustard crop, remove affected part and spray carbendazim @ 1 g/lit water in clear weather.
CHICK PEA	On the occurrence of fruit borer in chick pea crop, solution of 125 ml of Chlorantraniliprole 18.5 SL or 220 g of Emamectin Benzoate in 500-600 liter of water per hectare should be sprayed.
LENTIL	On the occurrence of fruit borer in lentil crop at the time of flowering, Melathiyan 50EC @ 2.0 liter/Ha or Quinalphos 25 EC @ 1.5liter/Ha in 500-600 liter of water should be sprayed.
BARLEY	On the occurrence of rust in crop, solution of Propiconazole 25 EC @ 1ml/liter of water should be sprayed by keeping in view the weather forecast.

### Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory		
ΤΟΜΑΤΟ	Drainage of excess water should be maintained in the field. If crop plants is showing fluffy cottony growth which leads to dry the plants. So infected plants must be removed and burn. In case of late blight, spray Mancozeb 64% + cymoxanil 8% WP @ 2.5gm per litre in clear weather by keeping in view the weather forecast.		
ροτατο	In case of late blight, spray Mancozeb $64\%$ + cymoxanil 8% WP @ 2.5gm p litre in clear weather by keeping in view the weather forecast.		
VEGETABLE PEA	On the occurrence of rust, triadimefon 25% WP @ 1 g/lit water should be sprayed.		

## Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
BUFFALO	Do not feed the animals with that green fodder which is in the early stage so that they do not get the disease called lactation tetany. To save the animals from cold, animal shed should be protected from cold and hot food should be given to the animals. To protect the animals from cold, Gur (Jaggery) with ajwain should be given to the animals without mixing in water. It gives warmth and energy to them.
COW	To save the animals from cold, animal shed should be protected from cold and hot food should be given to the animals. To protect the animals from cold, Gur (Jaggery) with ajwain should be given to the animals without mixing in water. It gives warmth and energy to them.

## **Poultry Specific Advisory:**

<b>Poultry(Varieties)</b>	Poultry Specific Advisory
HEN	There is a chance of Aflatoxicosis disease in Poultry due to occurrence of fungus in the food which causes to death of poultry birds. So, medicine should be administered to the poultry birds only on the advice of veterinarian.